

Ranking of Intentional self-harm as a leading Cause of Death

Idaho Residents 2003-2010

Year	Deaths		Ranking		Death Rates ¹			
	Number		Idaho	U.S.	Crude		Age Adjusted ²	
	of deaths	Percent			Idaho	U.S.	Idaho	U.S.
2003 ^{3,6}	218	-	9	-	16.0	10.5	16.3	10.5
2004 ⁷	239	2.4%	8	11	17.2	10.8	17.6	10.7
2005 ^{4,8}	225	2.1%	9	11	15.7	11.0	16.0	10.9
2006 ^{5,9}	218	2.1%	9	11	14.9	11.0	15.3	10.9
2007 ¹⁰	220	2.0%	9	11	14.7	11.1	14.9	10.9
2008 ¹⁰	251	2.3%	8		16.5	11.1	16.7	10.9
2009 ¹¹	307	2.8%	8		19.9	11.5	20.6	11.3
2010	290	2.5%	8		18.5		18.8	

Source: Idaho Vital Statistics 2003, Idaho Department of Health and Welfare, Division of Health, Bureau of Health Policy and Vital Statistics, 2005.
 Idaho Vital Statistics 2004, Idaho Department of Health and Welfare, Division of Health, Bureau of Health Policy and Vital Statistics, 2006.
 Idaho Vital Statistics 2005, Idaho Department of Health and Welfare, Division of Health, Bureau of Health Policy and Vital Statistics, 2007.
 Idaho Vital Statistics 2006, Idaho Department of Health and Welfare, Division of Health, Bureau of Vital Records and Health Statistics, 2008.

- Notes:
1. Rates are per 100,000 population and are based on tenth revision of the International Classification of Diseases (ICD-10)
 2. Age-adjusted rates are artificial measures developed to eliminate the bias inherent in differing age compositions, thus allowing estimate as the standard population comparisons between geographic regions. Idaho and U.S. age-adjusted rates were calculated using the 2000 U.S. population.
 3. The manner of coding the underlying cause of death changed in 1999 from the ninth revision (ICD-9) to the tenth revision of the comparability ratios based on Modified ICD-9 codes. International Classification of Diseases (ICD-10). Therefore, 2003 data are not comparable to data prior to 1999 without comparability ratios based on Modified ICD-9 codes.
 4. Idaho rates are based on 2005 population estimates based on 2000 Census.
 5. Idaho rates are based on July 1, 2006 population estimates based on 2000 Census.
 6. U.S. crude and age-adjusted rates are 2003 preliminary data. Rates are calculated using 2003 population estimates
 7. U.S. crude and age-adjusted rates are 2004 preliminary data. Rates are calculated using 2004 population estimates.
 8. U.S. crude and age-adjusted rates are 2004 final data for leading causes of death and are 2004 preliminary for sub-category
 9. U.S. crude and age-adjusted rates are 2005 final data for leading causes of death.
 10. U.S. crude and age-adjusted rates are 2006 final data for leading causes of death.
 11. U.S. crude and age-adjusted rates are 2007 data (latest final available) for leading causes of death.
 Rates are calculated using population estimates based on the 2000 Census.

